

The Weak $\Lambda\Lambda K^0$ Interaction

Martin J. Savage

Department of Physics, University of Washington, Seattle, WA 98195
savage@phys.washington.edu

Roxanne P. Springer

Duke University Department of Physics, Durham, NC 27708
rps@phy.duke.edu

Abstract

We calculate the leading contribution to the weak $\Lambda\Lambda K$ coupling in heavy baryon chiral perturbation theory, including the SU(3) breaking terms arising at one-loop. This coupling gives the leading one-boson exchange amplitude for the process $\Lambda\Lambda \rightarrow \Lambda n$ and is an important component in understanding the decay of $\Lambda\Lambda$ -hypernuclei. We find the S-wave and P-wave amplitudes to be stable under one-loop corrections, but the associated errors become large. We discuss the theoretical errors and potential effects of higher order terms.

I. INTRODUCTION

Experiments at Brookhaven (BNL-AGS-E885, E906) and the Japanese Hadron Facility (KEK-E373) are planned to measure the properties and decays of nuclei containing more than one strange baryon, such as $\Lambda\Lambda$ -hypernuclei. Some observations already exist [1]. In addition to the conventional decay mechanisms that exist for singly strange hypernuclei; mesonic decay such as $\Lambda \rightarrow N\pi$ (suppressed in heavy nuclei) and weak scattering such as $\Lambda N \rightarrow NN$, there is a decay amplitude for $\Lambda\Lambda$ -hypernuclei arising from $\Lambda\Lambda \rightarrow \Lambda N$. Unlike ΛN weak scattering there is no contribution from single π exchange to $\Lambda\Lambda$ weak scattering because of the absence of a strong $\Lambda\Lambda\pi$ coupling. The leading amplitude therefore comes from the exchange of a virtual K^0 . The strong ΛNK^0 interaction can be determined from the semi-leptonic decay $\Lambda \rightarrow pe^-\bar{\nu}_e$, using PCAC. However, the nonleptonic weak $\Lambda\Lambda K^0$ vertex cannot be measured directly. The approximate SU(3) flavor symmetry of the strong interactions can be used in combination with the assumption of octet-dominance in the nonleptonic weak interactions to relate the $\Lambda\Lambda K^0$ vertex to amplitudes for nonleptonic hyperon decays $B \rightarrow B'\pi$. In order to systematically include SU(3) breaking contributions to the decay we use chiral perturbation theory. Such calculations have been performed for the nonleptonic hyperon decays [2,3] and for the NNK vertices [4], assuming that the

breaking is dominated by terms of the form $m_s \log m_s$. We will determine the leading SU(3) breaking corrections of the form $m_s \log m_s$ to the $\Lambda\Lambda K^0$ couplings. These couplings may then be used as an input into the nuclear codes that presently exist (e.g., [5]) in order to determine the modification to the decay of $\Lambda\Lambda$ -hypernuclei.

An additional motivation for studying the $\Lambda\Lambda K^0$ couplings comes from an observation that some consider to cast doubt on the convergence of SU(3) chiral perturbation theory. A long-standing problem for chiral perturbation theory has been its failure to reliably predict the P-wave amplitudes in the nonleptonic decays of octet baryons. The loop corrections are relatively large compared to the tree level values [3] but this is thought to have more to do with the significant but accidental cancellation among tree level terms than an instability in the chiral expansion [6]. However, it is important to determine if the situation with the P-wave amplitudes in the octet decays is universal, and whether the predictions of chiral perturbation theory can be considered robust, with certain understood exceptions. To this end, we calculated the KNN couplings and found that the P-waves amplitudes are more stable against SU(3) breaking contributions than the S-waves amplitudes, and that neither is as badly behaved as P-wave amplitudes for octet baryon decay [4]. That work has been used [5] in an attempt to understand the discrepancy between the neutron induced versus proton induced decay rates of Λ hypernuclei. With the emergence of more accurate experimental results, we expect that the importance of the KNN coupling for these processes will be determined. We hope that the future experimental studies of $\Lambda\Lambda$ -hypernuclei and the substantial theoretical efforts presently underway will establish the $\Lambda\Lambda K^0$ vertex. Together with the data on singly-strange hypernuclei, the pattern of the weak octet baryon interactions will become clearer.

II. THE CHIRAL LAGRANGIAN FOR NONLEPTONIC INTERACTIONS

This section closely follows the section in [4] where the Lagrangian and relevant notation is presented. Weak nonleptonic hypernuclear decay takes place in an environment where typical momentum transfers are smaller than the chiral symmetry breaking scale; $p < \Lambda_\chi$, where $\Lambda_\chi \sim 1$ GeV. At these scales, it is appropriate to use as fundamental fields the lowest mass octet and decuplet baryons and the octet of pseudo-Goldstone bosons. The octet of baryons is

$$B_v = \begin{pmatrix} \frac{1}{\sqrt{2}}\Sigma_v^0 + \frac{1}{\sqrt{6}}\Lambda_v & \Sigma_v^+ & p_v \\ \Sigma_v^- & -\frac{1}{\sqrt{2}}\Sigma_v^0 + \frac{1}{\sqrt{6}}\Lambda_v & n_v \\ \Xi_v^- & \Xi_v^0 & -\frac{2}{\sqrt{6}}\Lambda_v \end{pmatrix}, \quad (2.1)$$

and the decuplet is

$$\begin{aligned} T_v^{111} &= \Delta_v^{++}, \quad T_v^{112} = \frac{1}{\sqrt{3}}\Delta_v^+, \quad T_v^{122} = \frac{1}{\sqrt{3}}\Delta_v^0, \quad T_v^{222} = \Delta_v^-, \\ T_v^{113} &= \frac{1}{\sqrt{3}}\Sigma_v^{*+}, \quad T_v^{123} = \frac{1}{\sqrt{6}}\Sigma_v^{*0}, \quad T_v^{223} = \frac{1}{\sqrt{3}}\Sigma_v^{*-}, \quad T_v^{133} = \frac{1}{\sqrt{3}}\Xi_v^{*0}, \\ T_v^{233} &= \frac{1}{\sqrt{3}}\Xi_v^{*-}, \quad T_v^{333} = \Omega_v^- \quad . \end{aligned} \quad (2.2)$$

The subscript v labels the four-velocity of the baryon. The average mass of the octet of baryons has been explicitly removed from the Lagrangian [8]. The octet of mesons is contained in

$$\Sigma = \xi^2 = \exp\left(\frac{2iM}{f}\right) \quad , \quad (2.3)$$

where

$$M = \begin{pmatrix} \frac{1}{\sqrt{6}}\eta + \frac{1}{\sqrt{2}}\pi^0 & \pi^+ & K^+ \\ \pi^- & \frac{1}{\sqrt{6}}\eta - \frac{1}{\sqrt{2}}\pi^0 & K^0 \\ K^- & \bar{K}^0 & -\frac{2}{\sqrt{6}}\eta \end{pmatrix} \quad . \quad (2.4)$$

Using this notation, a vector and an axial vector current can be defined:

$$\begin{aligned} V_\mu &= \frac{1}{2}(\xi\partial_\mu\xi^\dagger + \xi^\dagger\partial_\mu\xi) \\ A_\mu &= \frac{i}{2}(\xi\partial_\mu\xi^\dagger - \xi^\dagger\partial_\mu\xi) \quad . \end{aligned} \quad (2.5)$$

With these elements, and the covariant chiral derivative $\mathcal{D}_\mu = \partial_\mu + [V_\mu, \]$, an $SU(3)_L \otimes SU(3)_R$ symmetric Lagrangian can be formed. The lowest dimension operators will dominate observables, with the higher dimensions suppressed by increasing powers of $\frac{p}{\Lambda_\chi}$. To calculate the $\Lambda\Lambda K^0$ coupling to one loop, we need both strong and weak interactions of the baryons with the mesons,

$$\mathcal{L} = \mathcal{L}^{st} + \mathcal{L}^{wk} \quad . \quad (2.6)$$

The strong interactions are described by

$$\begin{aligned} \mathcal{L}^{st} &= i\text{Tr}\bar{B}_v(v \cdot \mathcal{D})B_v + 2D \text{Tr}\bar{B}_v S_v^\mu\{A_\mu, B_v\} + 2F \text{Tr}\bar{B}_v S_v^\mu[A_\mu, B_v] \\ &\quad - i\bar{T}_v^\mu(v \cdot \mathcal{D})T_{v\mu} + \Delta m \bar{T}_v^\mu T_{v\mu} + \mathcal{C}(\bar{T}_v^\mu A_\mu B_v + \bar{B}_v A_\mu T_v^\mu) \\ &\quad + 2\mathcal{H} \bar{T}_v^\mu S_{vv} A^\nu T_{v\mu} + \frac{f^2}{8} \text{Tr}\partial_\mu \Sigma \partial^\mu \Sigma^\dagger + \dots \quad , \end{aligned} \quad (2.7)$$

where f is the meson decay constant and only the lowest dimension operators are retained in the expansion. The axial couplings F, D, C , and $\mathcal{H}$ which appear in $\mathcal{L}^{(st)}$ could in principle be determined by matching this effective Lagrangian onto the appropriate calculation in QCD. Since such a computation in QCD has not been performed, these parameters are taken as unknowns and determined through a fit to experimental data. The $\Delta s = 1$ weak interactions required for this computation are described by the Lagrange density (assuming octet dominance)

$$\begin{aligned} \mathcal{L}^{wk} &= G_F m_\pi^2 f_\pi \left(h_D \text{Tr}\bar{B}_v\{\xi^\dagger h\xi, B_v\} + h_F \text{Tr}\bar{B}_v[\xi^\dagger h\xi, B_v] \right. \\ &\quad \left. + h_C \bar{T}_v^\mu(\xi^\dagger h\xi)T_{v\mu} + h_\pi \frac{f^2}{8} \text{Tr}(h\partial_\mu \Sigma \partial^\mu \Sigma^\dagger) + \dots \right) \quad , \end{aligned} \quad (2.8)$$

where

$$h = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad (2.9)$$

This Lagrange density contains additional unknown couplings that must be determined from data: h_D, h_F, h_π , and h_C . The pion decay constant is known to be $f_\pi \sim 132$ MeV. We have inserted factors of $G_F m_\pi^2 f_\pi$ in Eq. 2.8 so that the constants h_D, h_F , and h_C are dimensionless.

III. $\Lambda\Lambda K^0$ AMPLITUDES

The amplitude for the weak $\Delta s = 1$ nonleptonic interaction $\Lambda\Lambda K^0$ is given by

$$\mathcal{A} = iG_F m_\pi^2 \frac{f_\pi}{f_K} \bar{\Lambda}_v \left[\mathcal{A}^{(S)} + 2 \frac{k \cdot S_v}{\Lambda_\chi} \mathcal{A}^{(P)} \right] \Lambda_v, \quad (3.1)$$

where k is the outgoing momentum of the K^0 and

$$\mathcal{A}^{(L)} = \mathcal{A}_0^{(L)} + \mathcal{A}_1^{(L)} + \dots, \quad (3.2)$$

with $L=S$ for the S-wave amplitude, or $L=P$ for the P-wave amplitude. The subscripts in the above expansion for each L -wave indicate the order in chiral perturbation theory.

A. S-Wave Vertices

The tree level amplitude, as shown in Fig. 1 gives

$$\mathcal{A}_0^{(S)}(\Lambda\Lambda K^0) = -h_D. \quad (3.3)$$

The loop corrections are shown in Fig. 2 and evaluate to

$$\begin{aligned} \mathcal{A}_1^{(S)}(\Lambda\Lambda K^0) = & \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \left[\frac{10}{3} h_D + h_D \left(-\frac{11}{3} D^2 - 9F^2 \right) + 18h_F DF \right] \\ & + h_C C^2 \mathcal{J}(\Delta m) - \mathcal{A}_0^{(S)}(\Lambda\Lambda K^0) \mathcal{Z}_\Psi \end{aligned} \quad (3.4)$$

where the contribution from wavefunction renormalization (not shown in the figure) is given by

$$\mathcal{Z}_\Psi = \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \left(18F^2 + \frac{14}{3} D^2 \right) + C^2 \mathcal{J}(\Delta m), \quad (3.5)$$

and the function $\mathcal{J}(\delta)$ is

$$\mathcal{J}(\delta) = \frac{1}{16\pi^2 f_K^2} \left[(m_K^2 - 2\delta^2) \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) + 2\delta \sqrt{\delta^2 - m_K^2} \ln\left(\frac{\delta - \sqrt{\delta^2 - m_K^2 + i\epsilon}}{\delta + \sqrt{\delta^2 - m_K^2 + i\epsilon}}\right) \right]. \quad (3.6)$$

The argument of $\mathcal{J}$ is the mass splitting between the decuplet and octet of baryons. It is important to note that we have not included wavefunction renormalization for the K^0 so that double counting is avoided in the structure calculations [5].

B. P-Wave Vertices

The tree level P-wave amplitude arises from two pole graphs, as shown in Fig. 3 ,

$$\frac{\mathcal{A}_0^{(P)}(\Lambda\Lambda K^0)}{\Lambda_\chi} = \frac{1}{6} \frac{(D-3F)(3h_F - h_D)}{m_\Lambda - m_\Xi} - \frac{1}{6} \frac{(D+3F)(h_D + 3h_F)}{m_\Lambda - m_N} . \quad (3.7)$$

The loop level contributions are shown in Fig. 4 and Fig. 5. We present the contributions from each graph separately. As with the S-waves, the wavefunction renormalization on the external kaon is not included, since that will be accounted for in the structure calculations. We find that the full amplitude is the sum of the following contributions:

$$a = \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \left[\frac{D+3F}{m_\Lambda - m_N} \left(\frac{5}{12} h_D + \frac{5}{4} h_F \right) + \frac{D-3F}{m_\Lambda - m_\Xi} \left(\frac{5}{12} h_D - \frac{5}{4} h_F \right) \right] \quad (3.8)$$

$$b = \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \left[\frac{5}{36} \frac{(3F+D)(h_D + 3h_F)}{m_\Lambda - m_N} + \frac{5}{36} \frac{(3F-D)(3h_F - h_D)}{m_\Lambda - m_\Xi} \right] \quad (3.9)$$

$$c = \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \left[\frac{1}{6} \frac{(h_D + 3h_F)(D-3F)}{m_\Lambda - m_N} \left(\frac{19}{9} D^2 + \frac{4}{3} DF - 3F^2 \right) + \frac{1}{6} \frac{(3h_F - h_D)(D+3F)}{m_\Lambda - m_\Xi} \left(-\frac{19}{9} D^2 + \frac{4}{3} DF + 3F^2 \right) \right] \quad (3.10)$$

$$d = \mathcal{J}(\Delta m) C^2 \mathcal{H} \frac{5}{27} \frac{h_D + 3h_F}{m_\Lambda - m_N} + \mathcal{J}(\Delta m) C^2 \mathcal{H} \frac{5}{27} \frac{3h_F - h_D}{m_\Lambda - m_\Xi} \quad (3.11)$$

$$e = \frac{1}{9} \mathcal{K}(m_K, \Delta m) C^2 \frac{(h_D + 3h_F)(D-3F)}{m_\Lambda - m_N} - \mathcal{K}(m_K, \Delta m) C^2 \frac{h_D + 3h_F}{m_\Lambda - m_N} \frac{4D}{9} + \frac{3h_F - h_D}{m_\Lambda - m_\Xi} C^2 \left(-\frac{1}{3} (D-F) \mathcal{K}(m_K, \Delta m) - \frac{2}{9} D \mathcal{K}(m_\eta, \Delta m) \right) \quad (3.12)$$

$$f = \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \frac{(D+3F)(D-3F)}{m_\Lambda - m_N} \left(-\frac{2}{9} D(h_D + 3h_F) - \frac{1}{12} (D+3F)(h_D - 3h_F) - \frac{3}{4} (D-F)(h_D + h_F) \right) + \frac{m_K^2}{16\pi^2 f_K^2} \ln\left(\frac{m_K^2}{\Lambda_\chi^2}\right) \frac{(D+3F)(D-3F)}{m_\Lambda - m_\Xi} \left(-\frac{1}{12} (D-3F)(h_D + 3h_F) + \frac{2}{9} D(3h_F - h_D) + \frac{3}{4} (F+D)(h_F - h_D) \right) \quad (3.13)$$

$$g = \frac{1}{3} \mathcal{J}(\Delta m) C^2 h_C \frac{D + 3F}{m_\Lambda - m_N} + \frac{1}{3} \mathcal{J}(\Delta m) C^2 h_C \frac{3F - D}{m_\Lambda - m_\Xi} \quad (3.14)$$

$$h = -\frac{1}{36} D h_\pi \frac{m_K^2}{16\pi^2 f_K^2} \ln \left(\frac{m_K^2}{\Lambda_\chi^2} \right) \quad (3.15)$$

$$i = \frac{m_K^2}{16\pi^2 f_K^2} \ln \left(\frac{m_K^2}{\Lambda_\chi^2} \right) h_\pi D \left(\frac{7}{108} (D - 3F)^2 - \frac{1}{4} (D - F)(D + 3F) - \frac{1}{4} (F + D)(D - 3F) \right) \quad (3.16)$$

$$j = \mathcal{J}(\Delta m) C^2 \mathcal{H} h_\pi \frac{5}{18} \quad (3.17)$$

$$k = G_{m_\pi, m_K, \Delta m} C^2 h_\pi (D - 3F) \frac{1}{6} + G_{m_\pi, m_K, \Delta m} C^2 h_\pi \left(\frac{D}{6} - \frac{F}{2} \right) - G_{m_\eta, m_K, \Delta m} C^2 h_\pi \frac{D}{9} \quad (3.18)$$

$$\begin{aligned} l = & h_\pi (D - 3F) \left(\tilde{G}_{m_K, m_\eta, 0} \left(\frac{D^2}{16} - \frac{1}{8} D F - \frac{3}{16} F^2 \right) \right. \\ & - \tilde{G}_{m_K, m_\eta, \Delta m} \frac{1}{24} C^2 \\ & + \tilde{G}_{m_\pi, m_K, 0} \left(\frac{9}{16} F^2 - \frac{9}{8} F D - \frac{3}{16} D^2 \right) \\ & - \tilde{G}_{m_\pi, m_K, \Delta m} \frac{C^2}{4} \\ & + \frac{C^2 \Delta m}{m_\Lambda - m_\Xi} \left(\frac{1}{2} G_{m_\pi, m_K, \Delta m} + \frac{1}{12} G_{m_\eta, m_K, \Delta m} \right) \\ & + h_\pi (D + 3F) \left(\tilde{G}_{m_K, m_\eta, 0} \left(\frac{D^2}{16} + \frac{1}{8} F D - \frac{3}{16} F^2 \right) \right. \\ & + \tilde{G}_{m_\pi, m_K, 0} \left(-\frac{3}{16} D^2 + \frac{9}{8} F D + \frac{9}{16} F^2 \right) \\ & + \tilde{G}_{m_\pi, m_K, \Delta m} \frac{C^2}{8} \\ & \left. - \frac{1}{4} \frac{C^2 \Delta m}{m_\Lambda - m_N} G_{m_\pi, m_K, \Delta m} \right) \end{aligned} \quad (3.19)$$

$$\begin{aligned} \Psi_{off} = & -\frac{1}{6} \frac{(3h_F - h_D)(D - 3F)}{m_\Lambda - m_\Xi} \left[\frac{m_K^2}{16\pi^2 f_K^2} \ln \left(\frac{m_K^2}{\Lambda_\chi^2} \right) \left(\frac{17}{3} D^2 + 10 D F + 15 F^2 \right) + \frac{13}{3} C^2 \mathcal{J}(\Delta m) \right] \\ & + \frac{1}{6} \frac{(h_D + 3h_F)(D + 3F)}{m_\Lambda - m_N} \left[\frac{m_K^2}{16\pi^2 f_K^2} \ln \left(\frac{m_K^2}{\Lambda_\chi^2} \right) \left(\frac{17}{3} D^2 - 10 D F + 15 F^2 \right) + C^2 \mathcal{J}(\Delta m) \right] \end{aligned} \quad (3.20)$$

$$\Psi = -\mathcal{A}_0^{(P)}(\Lambda\Lambda K^0)\mathcal{Z}_\Psi \quad (3.21)$$

The external baryon wavefunction renormalization is contained in Ψ , with the function $\mathcal{Z}_\Psi$ defined as before in the S-wave formulas. The additional functions which appear in the above expressions are given by

$$\begin{aligned} \mathcal{K}(m, \delta) = \frac{1}{16\pi^2 f_K^2} & \left\{ (m^2 - \frac{2}{3}\delta^2) \ln\left(\frac{m^2}{\Lambda_\chi^2}\right) \right. \\ & \left. + \frac{2}{3} \frac{1}{\delta} \left[(\delta^2 - m^2)^{3/2} \ln\left(\frac{\delta - \sqrt{\delta^2 - m^2 + i\epsilon}}{\delta + \sqrt{\delta^2 - m^2 + i\epsilon}}\right) + \pi m^3 \right] \right\} \quad , \end{aligned} \quad (3.22)$$

$$\begin{aligned} G_{m_1, m_2, \delta} &= \frac{m_1^2}{m_1^2 - m_2^2} \mathcal{K}(m_1, \delta) + \frac{m_2^2}{m_2^2 - m_1^2} \mathcal{K}(m_2, \delta) \quad , \\ \tilde{G}_{m_1, m_2, \delta} &= \frac{m_1^2}{m_1^2 - m_2^2} \mathcal{J}(m_1, \delta) + \frac{m_2^2}{m_2^2 - m_1^2} \mathcal{J}(m_2, \delta) \quad . \end{aligned} \quad (3.23)$$

As before, the argument δ is the average decuplet-octet mass splitting.

IV. NUMERICAL RESULTS

Our numerical results for the $\Lambda\Lambda K^0$ couplings depend upon a number of parameters that must be fit to experimental data. For tree level estimates, we use parameters obtained by fitting tree level expressions to the experimentally measured rates and asymmetries in the semileptonic decays of octet baryons, and the S-wave amplitudes in the nonleptonic decay of octet baryons. The systematic theoretical error associated with the neglect of SU(3) breaking is included in the fit by increasing the error bars of the experimental measurements to 20%. We find that the tree level expressions for the S- and P- wave amplitudes are

$$\begin{aligned} \mathcal{A}_0^{(S)}(\Lambda\Lambda K^0) &= 0.55 \pm 0.19 \\ \frac{\mathcal{A}_0^{(P)}(\Lambda\Lambda K^0)}{\Lambda_\chi} &= -5.2 \pm 1.0 \quad , \end{aligned} \quad (4.1)$$

using the fitted values $D = 0.82 \pm 0.07$, $F = 0.48 \pm 0.06$, $h_D = -0.55 \pm 0.19$, and $h_F = 1.37 \pm 0.11$.

Loop level numerical results for the $\Lambda\Lambda K^0$ couplings depend upon parameters extracted at loop level from other processes. We find that simultaneously fitting the semileptonic decays of the octet of baryons [9], the strong decays of the decuplet of baryons [10], and the S-wave nonleptonic decays of the octet of baryons [3] yields for $\Delta m = 0$

$$\begin{aligned} \mathcal{A}_0^{(S)}(\Lambda\Lambda K^0) + \mathcal{A}_1^{(S)}(\Lambda\Lambda K^0) &= 0.43 \pm 0.89 \\ \mathcal{A}_0^{(P)}(\Lambda\Lambda K^0) + \mathcal{A}_1^{(P)}(\Lambda\Lambda K^0) &= -4.9 \pm 1.9 \quad . \end{aligned} \quad (4.2)$$

This suggests that the S-wave predictions are probably unreliable in chiral perturbation theory. When $\Delta m = 200$ MeV is used for the $\Lambda\Lambda K^0$ couplings, we obtain

$$\begin{aligned}
\mathcal{A}_0^{(S)}(\Lambda\Lambda K^0) + \mathcal{A}_1^{(S)}(\Lambda\Lambda K^0) &= 0.28 \pm 0.82 \\
\mathcal{A}_0^{(P)}(\Lambda\Lambda K^0) + \mathcal{A}_1^{(P)}(\Lambda\Lambda K^0) &= -4.8 \pm 4.1 \quad .
\end{aligned}
\tag{4.3}$$

Both the S-wave and P-wave amplitudes have very large uncertainties associated with them. The S-wave amplitude is small at tree-level and it is no surprise to find the loop corrections substantially modify the leading result. However, for the P-waves the tree-level amplitude is not small and it would be somewhat disturbing to find large cancellations between the higher order and tree level amplitudes. A significant cancellation may suggest a failure of chiral perturbation theory.

In the fit we find [11,12]¹ $|C| = 1.11 \pm 0.05$, $D = 0.54 \pm 0.03$, $F = 0.33 \pm 0.03$, $\mathcal{H} = -1.75 \pm 0.46$, $h_C = 0.30 \pm 6.75$, $h_D = -0.42 \pm 0.21$, and $h_F = 0.94 \pm 0.44$ when a decuplet-octet mass splitting of zero is used. The sign of C is not determined as it can be changed by a field redefinition and presents no ambiguity in our results. The errors on h_C are large because the nonleptonic decays of octet baryons depend on h_C only through loops. The difference between the tree level and loop level extracted parameters is larger than one might expect from naive dimensional analysis. However, it is only observables that are required to be stable in order to have confidence that the chiral expansion is converging.

V. DISCUSSION

The S-wave and P-wave amplitudes for the $\Lambda\Lambda K^0$ interaction are explored at loop level in chiral perturbation theory. This adds to the body of knowledge on $\Delta s = 1$ transitions in chiral perturbation theory already existing from the nonleptonic hyperon decays, and the KNN couplings which can be measured in single Λ hypernuclear decay. Comparing these calculations with data yields insight into the strengths and weaknesses of the theory, and highlight which results can be considered typical for the theory, and which anomalous.

The tree-level $\Lambda\Lambda K$ S-wave amplitude is small, making it vulnerable to higher order corrections. The S-wave amplitudes for both nonleptonic hyperon decays and the KNN couplings are generally of order one or larger at tree level. The tree-level $\Lambda\Lambda K$ P-wave amplitude, on the other hand, is substantial. Like the P-wave nonleptonic hyperon decay amplitude, the $\Lambda\Lambda K$ coupling has contributions from two pole diagrams. But unlike the nonleptonic hyperon amplitudes, a significant cancellation does not occur. This lends credence to the claim that the situation with the nonleptonic hyperons may be anomalous. For the KNN amplitudes, only one P-wave diagram contributes and no cancellation is possible. The errors associated with these results come from the propagation of errors associated with fitting other observables at tree-level and are of the expected magnitude.

Looking at the one-loop results with $\Delta m = 0$, we note that the central values are not dramatically different from the tree level values, but the S-wave errors have increased to the point where we question whether the chiral expansion is converging for this coupling. This is in contrast to the situation with the nonleptonic hyperon decays where the experimental S-waves are well reproduced at loop level. The KNN S-wave couplings suffer from fairly

¹ We cannot reproduce the values of axial couplings (or uncertainties) found in [13].

large (up to 50%) corrections from one loop diagrams, but the associated errors are small enough that the predictions remain meaningful. The P-waves for the nonleptonic hyperon decays poorly reproduce the data at one loop and experience large corrections. If it is true that this is caused by the accidental cancellations that make the tree level values small, we should not see a similar trend for the KNN or $\Lambda\Lambda K$ couplings. Indeed, the KNN P-wave loop corrections are not large, and the associated errors are reasonable. The predictions for $\Lambda\Lambda K$ with $\Delta m = 0$ also look very stable, with a small correction and small errors. This alone might make us think that the P-wave couplings are generally well behaved but that predictions for processes which are anomalously small at tree level cannot be trusted. Once again, the errors come from a propagation of the errors from the global parameter fit. They are stable under reasonable changes in the meson decay constant, and the chiral symmetry breaking scale.

An idea of the stability of our result to higher order corrections can be obtained by considering the situation when $\Delta m \neq 0$. The P-wave amplitude acquires a significantly increased error. It is interesting that, despite this, the central value remains basically unchanged. Yet the large error indicates that the P-wave $\Lambda\Lambda K$ coupling could be sensitive to higher order corrections. In contrast to the P-wave amplitudes for the nonleptonic decay of octet baryons, where the trouble comes from accidental cancellations between pole graphs at tree level, here the difficulty comes because the parameters appear in the $\Lambda\Lambda K$ formula with nonzero Δm in such a way that errors on the parameters conspire to make the error on the result large. The S-wave amplitude in $\Lambda\Lambda K^0$ suffers both from a large uncertainty and a small tree level value. This situation is very different from that for the NNK amplitudes, where the uncertainties even with nonzero Δm were found to be reasonably small, $\sim 20\%$ or smaller.

In summary, we find that at one-loop level (without local counterterms which begin at $\mathcal{O}(m_s)$, formally higher order) the uncertainty in the $\Lambda\Lambda K$ amplitudes appears to nearly 100% for the P-waves and more than 200% for the S-waves. These numbers include an estimate of the theoretical error resulting from the neglect of higher order terms in the chiral expansion. The indication is that the $\Lambda\Lambda K$ coupling may be one of the quantities which is not well predicted by a loop level calculation in chiral perturbation theory. This is not the case for other processes. The fact that the parameters in one loop calculations fit the known data on axial currents, the strong decays of the decuplet, and the S-wave nonleptonic hyperon decays so well shows that, for these processes, higher order counter terms are not required.

A measurement of the P-wave $\Lambda\Lambda K$ coupling through the decay of $\Lambda\Lambda$ hypernuclei will yield important information on chiral perturbation theory, its parameters, and the importance of higher order terms. We are encouraged by the promise of experiments at KEK and BNL, and hope that possibilities at Frascati and TJNAF will also be pursued.

This was supported in part by a grant from the Department of Energy, DE-FG02-96ER40945. We thank C. Bennhold for several discussions that led to this work. We also thank T. Fakuda, E. Hungerford, and I. Melnikov for helpful communications.




FIG. 1. Tree level diagram for the $\Lambda\Lambda K^0$ amplitude. The black dot indicates the weak vertex.

REFERENCES

- [1] M. Danysz, et al., Phys. Rev. Lett. **11**, 29 (1993); D.J. Prowse, Phys. Rev. Lett. **17**, 782 (1966); S. Aoki et al., Prog. Theor. Phys. **89**, 493 (1993).
- [2] J. Bijnens, H. Sonoda and M.B. Wise, Nucl. Phys. **B261**, 185 (1985).
- [3] E. Jenkins, Nucl. Phys. **B375** 561 (1992).
- [4] M.J. Savage and R.P. Springer, Phys. Rev. C **53**, 441 (1996).
- [5] A. Parreno, A. Ramos, and C. Bennhold, Phys. Rev. C **56**, 339 (1997).
- [6] C. Carone and H. Georgi, Nucl. Phys. **B375** 243 (1992).
- [7] A. Ramos, E. van Meijgaard, C. Bennhold, and B.K. Jennings, Nucl. Phys. **A544**, 703 (1992).
- [8] The formalism of heavy baryon chiral perturbation theory is introduced in E. Jenkins and A.V. Manohar, In “Dobogokoe 1991, Proceedings, *Effective field theories of the standard model*” 113-137.
- [9] E. Jenkins and A. Manohar, Phys. Lett. **B255**, 558 (1991); **B259**, 353 (1991).
- [10] M.N. Butler, M.J. Savage and R.P. Springer, Nucl. Phys. **B399**, 69 (1993).
- [11] R.P. Springer, DUKE-TH-97-152.
- [12] M.J. Savage and J. Walden, Phys. Rev. D **55** 5367 (1997).
- [13] M.A. Luty and M. White, Phys. Lett. **B319**, 261 (1993).




FIG. 2. Loop level diagrams for the S-wave amplitude. The dashed lines are mesons, the solid lines are octet baryons, and the double lines are decuplet baryons. An unmarked vertex represents a strong interaction and the black dots are weak interactions.




FIG. 3. Tree level diagrams for the $\Lambda\Lambda K^0$ amplitude. An unmarked vertex represents a strong interaction and the black dots are weak vertices.




FIG. 4. Loop level diagrams for the P-wave amplitude. The dashed lines are mesons, the solid lines are octet baryons, and the double lines are decuplet baryons. An unmarked vertex represents a strong interaction and the black dots are weak interactions.




FIG. 5. Continuation of loop level diagrams for the P-wave amplitude. The dashed lines are mesons, the solid lines are octet baryons, and the double lines are decuplet baryons. An unmarked vertex represents a strong interaction and the black dots are weak interactions.